

| | | |
|-----------|---|-----|
| Figure 25 | Downhole geophysical log from petroleum well Mondarra 4 (1200–2200 m bgl) showing the Cattamarra Coal Measures and Eneabba Formation..... | 86 |
| Figure 26 | Downhole geophysical log from petroleum well Eneabba 1 (1700–3000 m bgl) showing the Cattamarra Coal Measures and Eneabba Formation..... | 87 |
| Figure 27 | Downhole geophysical log of petroleum well Ocean Hill 1 (2700–3700 m bgl) showing the Cadda Formation | 92 |
| Figure 28 | Downhole geophysical log of petroleum well Warro 1 (600–4385 m bgl) showing the Yarragadee Formation | 96 |
| Figure 29 | Downhole geophysical log of petroleum well Ocean Hill 1 (0–3000 m bgl) showing the Yarragadee Formation | 97 |
| Figure 30 | Yarragadee Formation Unit A: contours on base of unit..... | 100 |
| Figure 31 | Yarragadee Formation Unit B: contours on base of unit..... | 101 |
| Figure 32 | Yarragadee Formation Unit C: contours on base of unit..... | 102 |
| Figure 33 | Yarragadee Formation Unit D: contours on base of unit..... | 103 |
| Figure 34 | Downhole geophysical log from Gillingarra Line GL6 (0–975 m bgl) showing the Parmelia Group..... | 107 |
| Figure 35 | Parmelia Group: contours on base of unit..... | 108 |
| Figure 36 | Downhole geophysical log from Gillingarra Line 2B (0–1000 m bgl) showing the Warnbro Group | 114 |
| Figure 37 | Warnbro Group: contours on base of unit | 115 |
| Figure 38 | Gage Sandstone: contours on base of unit | 116 |
| Figure 39 | South Perth Shale: contours on base of unit | 117 |
| Figure 40 | Leederville Formation: contours on base of unit..... | 118 |
| Figure 41 | Downhole geophysical log from Gillingarra Line GL8 (0–1170 m bgl) showing the Coolyena Group..... | 127 |
| Figure 42 | Coolyena Group: contours on base of unit..... | 128 |
| Figure 43 | Henley Sandstone Member: contours on base of unit | 129 |
| Figure 44 | Kardinya Shale Member: contours on base of unit..... | 130 |
| Figure 45 | Mirrabooka Member: contours on base of unit | 131 |
| Figure 46 | Molecap Greensand (including Gingin Chalk and Poison Hill Greensand) and Lancelin Formation: contours on base of unit..... | 132 |
| Figure 47 | Surficial deposits: extent | 137 |
| Figure 48 | Digital elevation mapping showing Yallalie Basin..... | 138 |
| Figure 49 | Superficial formations: surface geology with geological cross-sections..... | 141 |
| Figure 50 | Superficial formations: contours on base of unit..... | 142 |
| Figure 51 | Geological cross-sections showing stratigraphical relationships of superficial formations..... | 143 |
| Figure 52 | Ascot Formation: isopachs..... | 146 |
| Figure 53 | Aquifers and aquitards below the superficial formations or surficial deposits | 160 |
| Figure 54 | Bore sites with aquifer test data (bores in the Guilderton–Lancelin area are within the Superficial aquifer) | 161 |
| Figure 55 | Hydrogeological cross-sections (hydraulic head) | 162 |
| Figure 56 | Hydrogeological cross-sections (salinity) | 165 |
| Figure 57 | Hydrogeological setting across the Northampton Inlier | 168 |
| Figure 58 | Northern region: hydrogeology and potentiometric surface | 169 |
| Figure 59 | Northern region: groundwater salinity..... | 170 |
| Figure 60 | Representative depth to watertable (2015) | 171 |
| Figure 61 | Water level change between mid-1980s and mid-2000s | 172 |